WISCONSIN

CANCER BURDEN



In 2004, the American Cancer Society estimates that¹

- 1,368,030 new cancer cases will be diagnosed in the United States, including 26,160 in Wisconsin.
- 563,700 cancer deaths will occur in the United States, including 10,780 in Wisconsin.

The average annual age-adjusted death rate for cancer per 100,000 persons²

Wisconsin: 194.3 National: 199.8

LUNG CANCER

In 2004, the American Cancer Society estimates that¹

- 3,050 new cases of lung cancer will be diagnosed among men and women in Wisconsin.
- 2,820 men and women will die of lung cancer in Wisconsin.

The CDC National Center for Health Statistics provides the following death rates:

The average annual age-adjusted death rates for lung cancer per 100,000 persons, by race, 1997–2001^{2,3}

| | Wisconsin | National |
|------------------------------|-----------|----------|
| Overall | 49.5 | 56.2 |
| White | 48.8 | 56.2 |
| Black | 76.9 | 65.2 |
| Hispanic | 14.5 | 24.9 |
| Asian/Pacific Islander | 25.5 | 28.2 |
| American Indian/Alaska Nativ | e 64.2 | 36.3 |

BREAST CANCER

In 2004, the American Cancer Society estimates that¹

- 4,040 new cases of breast cancer will be diagnosed among women in Wisconsin.
- 750 women will die of breast cancer in Wisconsin.

The CDC National Center for Health Statistics provides the following death rates:

The average annual age-adjusted death rates for breast cancer per 100,000 persons, by race, 1997–2001^{2,3}

| | Wisconsin | National |
|------------------------------|-----------|----------|
| Overall | 26.2 | 27.0 |
| White | 26.1 | 26.4 |
| Black | 31.6 | 35.4 |
| Hispanic | - | 17.2 |
| Asian/Pacific Islander | - | 12.6 |
| American Indian/Alaska Nativ | e - | 13.6 |

Visit http://www.cdc.gov/cancer/npcr/index.htm — *United States Cancer Statistics* — for more information on top cancer sites by geographic area, race, and gender.

COLORECTAL CANCER

In 2004, the American Cancer Society estimates that¹

- 2,900 new cases of colorectal cancer will be diagnosed among men and women in Wisconsin.
- 1,120 men and women will die of colorectal cancer in Wisconsin.

The CDC National Center for Health Statistics provides the following death rates:

The average annual age-adjusted death rates for colorectal cancer per 100,000 persons, by race, 1997–2001^{2,3}

| | Wisconsin | National |
|-------------------------------|-----------|----------|
| Overall | 20.7 | 20.8 |
| White | 20.5 | 20.3 |
| Black | 27.9 | 28.3 |
| Hispanic | - | 14.2 |
| Asian/Pacific Islander | - | 13.0 |
| American Indian/Alaska Nativo | e 29.9 | 13.9 |

PROSTATE CANCER

In 2004, the American Cancer Society estimates that¹

- 3,850 new cases of prostate cancer will be diagnosed among men in Wisconsin.
- 500 men will die of prostate cancer in Wisconsin.

The CDC National Center for Health Statistics provides the following death rates:

The average annual age-adjusted death rates for prostate cancer per 100,000 men, by race, 1997–2001^{2,3}

| | Wisconsin | National |
|------------------------------|-----------|----------|
| Overall | 32.6 | 31.5 |
| White | 32.1 | 28.8 |
| Black | 62.4 | 70.4 |
| Hispanic | - | 23.6 |
| Asian/Pacific Islander | - | 13.0 |
| American Indian/Alaska Nativ | e - | 20.2 |

¹ Source: American Cancer Society Facts and Figures, 2004. Estimates exclude more than a million cases of basal and squamous cell skin cancers and *in situ* cancers, except urinary bladder, that will be diagnosed in 2004. Lung cancer rates include bronchus cancer. State death totals were rounded to nearest 10.

² Source: Centers for Disease Control and Prevention (CDC) National Center for Health Statistics, vital statistics data, underlying cause of death, 1997–2001. Death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population.

³ Hyphens represent suppression of rates when there were 75,000 or fewer persons in the denominator or 20 or fewer deaths in the numerator.